



REC'D PCT/PTO 16 JUN 2005 #3

PCT/GB 2003 / 0 8 5 4 5 0

U 9 DECEMBER 2003

INVESTOR IN PEOPLE

## PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

REC'D 13 JAN 2004  
WIPO PCT


The Patent Office  
Concept House  
Cardiff Road  
Newport  
South Wales  
NP10 8QQ

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



*L. Mahoney*

Signed

Dated 8 October 2003

### BEST AVAILABLE COPY



24DEC02 EPO2875-1 D02693  
P01/7700 0.00-0229803.2

# Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)



The Patent Office

Cardiff Road  
Newport  
South Wales  
NP10 8QQ

1. Your reference PPD 70182/GB/P

2. Patent application number  
(The Patent Office will fill in this part) 20 DEC 2002 0229803.2

3. Full name, address and postcode of the or of each applicant (underline all surnames)  
SYNGENTA Limited  
European Regional Centre  
Priestley Road  
Surrey Research Park, Guildford,  
Surrey, GU2 7YH, United Kingdom  
  
Patents ADP number (if you know it) 06244007002  
If the applicant is a corporate body, give the country/state of its incorporation UNITED KINGDOM 0833 0748 001

4. Title of the invention CHEMICAL PROCESS

5. Name of your agent (if you have one) John Richard WATERMAN  
Intellectual Property Department  
Syngenta Limited  
Jealott's Hill International Research Centre  
PO Box 3538  
Bracknell, Berkshire, RG42 6YA  
UNITED KINGDOM  
  
"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

Patents ADP number (if you know it) 06791534002

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number	Country	Priority application number (if you know it)	Date of filing (day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application	Number of earlier application	Date of filing (day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

- a) any applicant named in part 3 is not an inventor, or
- b) there is an inventor who is not named as an applicant, or
- c) any named applicant is a corporate body.

See note (d) YES (b)

**Patents Form 1/77**

9. Enter the number of sheets for of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description

07

Claim(s)

01

Abstract

01

Drawing(s)

00

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patents Form 9/77)

Request for substantive examination (Patents Form 10/77)

Any other documents  
(please specify)

11.

I/We request the grant of a patent on the basis of this application.  
Syngenta Limited

Signature M A Rudd  
Authorised Signatory

Date 20/12/2002

12. Name and daytime telephone number of person to contact in the United Kingdom

Margaret Ann RUDD = 01344 413673

**Warning**

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

**Notes**

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 08459 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.


# CHEMICAL PROCESS

The present invention relates to a process for making insecticidal cyclopropanecarboxylic acid esters. More particularly, the invention relates to a process for making gamma-cyhalothrin [(S)- $\alpha$ -cyano-3-phenoxybenzyl (Z)-(1R,3R)-3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethylcyclopropanecarboxylate].

It is well known that the insecticidal activity of pyrethroids such as cyclopropane-carboxylic acid esters e.g. cyhalothrin is greatly affected by their stereochemistry. It is disclosed in Bentley et al, Pestic.Sci. (1980), 11(2), 156-64) that (S)- $\alpha$ -cyano-3-phenoxybenzyl (Z)-(1R,3R)-3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethylcyclopropanecarboxylate is the most active isomer of cyhalothrin.


Known processes for making gamma-cyhalothrin, such as those outlined in GB2000764, EP132392, involve esterification of 1R *cis*-Z 3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethylcyclopropanecarboxylic acid with an optically active cyanohydrin. However on an industrial scale it is preferable to find alternative methods of making the final product that avoid the use of the very expensive cyanohydrin.

There is therefore provided a process for the preparation of gamma-cyhalothrin comprising a) chlorinating 1R *cis*-Z 3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethylcyclopropanecarboxylic acid (V) to give 1R *cis*-Z 3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethylcyclopropanecarboxylic acid chloride (III); b) esterifying 1R *cis*-Z 3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethylcyclopropanecarboxylic acid chloride (III) with 3-phenoxybenzaldehyde (IV) in the presence of a source of cyanide to form a diastereoisomeric mixture of 1R cyhalothrin isomers (II) and c) epimerising the diastereoisomeric mixture (II) under conditions in which the least soluble diastereoisomer (I) crystallises from solution.



(V)

(III)



1R *cis*-Z 3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethyl cyclopropanecarboxylic acid (V) is a known compound and its preparation is described for example in US4683089, WO02/06202, WO97/03941 and WO/9942432.

Step a) is performed by standard techniques as in '*March 4<sup>th</sup> Edition - p437-38*'.

Preferred chlorinating agents are thionyl chloride, phosgene or phosphorous oxychloride.

It is preferred that solvents used in the process are hydrocarbons such as toluene, hexane, heptane or fluorobenzene.

Preferred temperatures for the process are from ambient to 100°C or the boiling point of the solvent

Preferably the acid (V) has an enantiomeric purity of greater than 80% of 1R 3R enantiomer, and most preferably greater than 90% 1R 3R enantiomer.

Step b) is performed in the presence of a cyanide source. Suitable sources of cyanide are alkali or alkaline earth metal cyanides, but preferably an alkali metal cyanide and more preferably sodium cyanide.

The reaction can be carried out in a solvent or in a mixture of a water immiscible solvent and an aqueous solution of the source of cyanide or in the absence of a solvent. Suitable solvents are aliphatic or aromatic hydrocarbons solvents. Examples of aromatic hydrocarbons are toluene, o-xylene, mixed xylenes or halobenzenes, for example fluorobenzene. Aliphatic hydrocarbons are for example hexane, iso-hexane, heptane, octane or mixtures of hydrocarbons commonly known as petroleum ethers. Preferred solvents are hexane, iso-hexane, heptane or octane.

The acid chloride (III) and 3-phenoxybenzaldehyde (IV) can be added sequentially or simultaneously to the source of cyanide in the presence of the solvent or the solvent mixture, optionally in presence of an organic base or an onium salt.

Alternatively, the source of cyanide may be added to a mixture of the acid chloride (III) and 3-phenoxybenzaldehyde in the presence of a solvent, optionally in the presence of water, if a water-immiscible solvent is used, and optionally in the presence of an organic base or an onium salt.

5 Suitably, the source of cyanide is used in excess, preferably between 1.0 and 2.0 molar equivalents on the acid chloride and most preferably 1-1.5 equivalents.

The aldehyde (IV) is used in substantially stoichiometric amounts, but may be used in a slight excess of up to 0.10 equivalents on the acid chloride.

10 The organic base is preferably a tertiary amine, such as triethylamine, triisopropylamine, or a tertiary diamine such as tetramethylethylene diamine, 1,4-diazabicyclo[2,2,2]octane or 1,4-diazabicyclo[5.4.0]undec-7-ene. The base can be used in sub-stoichiometric amounts, preferably 0-50mol% on the acid chloride and most preferably 1-10mol%. The onium salt can be a quaternary ammonium salt, such as tetra n-butylammonium bromide, or a phosphonium salt or a sulphonium or sulfoxonium salt, and is used in similar proportions as the organic base.

15 The reaction can be carried out at between -10°C and the boiling point of the solvent system, preferably between 0°C and 60°C and most preferably between 0°C and 20°C.

The compound of formula (II) may be isolated and purified or used directly in step c) without isolation or purification.

20 The compound of formula (II) is a mixture of (S)- $\alpha$ -cyano-3-phenoxybenzyl (Z)-(1R,3R)-3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethylcyclopropanecarboxylate and (R)- $\alpha$ -cyano-3-phenoxybenzyl (Z)-(1R,3R)-3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethylcyclopropanecarboxylate and is novel and as such it forms a further aspect of the invention.

Compound (IV), 3-phenoxy benzaldehyde, is a commercially available compound

25 Step c) may also be carried out in a solvent and a source of cyanide. The solvent and the source of cyanide are preferably those used in step b) although the cyanide need only be present in catalytic amounts, typically from 1-10mol%. The organic base optionally present in step b) may also be optionally present. Step c) may be carried out in the presence of a seed of gamma-cyhalothrin where the seed is present in an amount of between 1% and 90% of the amount of the 30 1R cyhalothrin entering step c). The seed is preferably dispersed in the solvent before cooling at the start of step c), or may be the retained crystallization mass from a previous execution of step c).

The reaction is performed at  $-20^{\circ}\text{C}$  to  $50^{\circ}\text{C}$ , preferable  $-20^{\circ}\text{C}$  to  $30^{\circ}\text{C}$  and most preferably  $-10^{\circ}\text{C}$  to  $10^{\circ}\text{C}$ . On completion of the reaction, the slurry can be filtered to recover the gamma-cyhalothrin, preferably having first added an acid to arrest the epimerisation reaction. The acid can be a mineral acid such as hydrochloric acid or sulphuric acid or an organic acid such as formic or acetic acid and is used in at least stoichiometric amounts on the total bases present in the system, i.e. the organic base plus any residual source of cyanide, and preferably in twice stoichiometric amounts. The solid gamma-cyhalothrin can be washed and dried by conventional methods. Alternatively, the slurry of gamma-cyhalothrin in the reaction mixture after addition of acid can be heated to dissolve the product, washed with water to remove catalysts and salts and the aqueous phase removed by separation. As appreciated by those skilled in the art, the pH of any water washes can be adjusted to facilitate removal of acid or basic catalysts or by-product species added or formed during the process. The gamma-cyhalothrin can then be recovered from the solvent layer by, for example, cooling, crystallization and filtration, or by evaporation of the solvent.

Surprisingly, the applicants have also found it is possible to make gamma-cyhalothrin in a process wherein steps b) and c) can be combined (in a one-pot process), that is a process for the preparation of gamma-cyhalothrin in which 1R *cis*-Z 3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethyl cyclopropanecarboxylic acid chloride (III) is reacted with 3-phenoxy benzaldehyde (IV) and a source of cyanide to give gamma-cyhalothrin (I).

The reaction is performed in a single step and does not require the isolation of the mixture of diastereomers (II) formed in the initial esterification. This embodiment has the advantage of reducing solvent recovery/waste disposal, of enabling higher productivity with less equipment and of reducing impurity formation due to the avoidance of reactive solvents and reduced work-up times.

It is preferred to carry out the reaction in a solvent.

In the combined process, once the esterification is complete, as judged by analysis, it is advantageous to remove any aqueous phase, after adjusting the pH to between pH 9-13, by warming the reaction mixture to  $20^{\circ}\text{C}$ - $60^{\circ}\text{C}$  and separation, before substantial amounts of epimerisation and crystallization of gamma-cyhalothrin takes place. The pH can be adjusted with addition of an inorganic base such as sodium hydroxide or sodium carbonate. The reaction mixture can then be cooled to the preferred epimerisation and crystallization temperature to complete the process. Optionally the organic base, or an onium salt and additional source of

cyanide can be added after the esterification, pH adjustment and separation but before substantial amounts of epimerisation and crystallization of gamma-cyhalothrin has taken place. In this situation, the cyanide need only be added in catalytic amounts, typically from 1-10mol%.

The following Examples illustrate the invention.

- 5 The products were analysed by Gas Chromatography using an Agilent gas chromatograph with a Chrompack CP Sil 5 CB column (50 metres, 0.32 mm ID and 0.1  $\mu$ m film thickness) with helium as carrier, split injection at 15 psi. Injection temperature 300°C detector 325°C and a detector gas composition of hydrogen 30 ml/min, air 350 ml/min and helium at 30 ml/min). The oven temperature profile was: initial temp 50°C, initial time 6 mins then heating rate 10°C/min to 120°C and hold for 3 mins then ramp to 240°C at 25°C/min. Hold for 8 minutes then ramp to 300°C at 50°C and hold for 6 minutes to burn off the column.

Using these conditions, the following retention times were observed:

- (S)- $\alpha$ -cyano-3-phenoxybenzyl (Z)-(1R,3R)-3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethylcyclopropanecarboxylate (gamma-cyhalothrin) 27.4 mins
- 15 (R)- $\alpha$ -cyano-3-phenoxybenzyl (Z)-(1R,3R)-3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethylcyclopropanecarboxylate 27.0 mins

#### EXAMPLE 1

##### **Preparation of 1R *cis*-Z 3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethyl-cyclopropane carboxylic acid chloride**

- 20 A 1 litre dry, clean jacketed split reaction vessel equipped with agitator, thermometer, condenser, nitrogen blanket and vent to a scrubber system was charged with toluene (450ml) and agitated whilst 1R *cis*-Z 3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethyl-cyclopropane carboxylic acid (89.4gm = 0.369mol) was added followed by triethylamine (0.21gm = 2.1mmol). The reaction mixture was then heated to 45°C, using oil circulation on the jacket, and thionyl chloride (62.0gm = 0.52mol) was then charged over 105 minutes maintaining on temperature. The reaction mass was then agitated for 5 hours at 45°C then tested by GLC for completion of reaction showing 2% residual acid. A further addition of thionyl chloride (4.4gm = 37mmol) was then made and the reaction mass allowed to cool with stirring overnight. The following day, residual thionyl chloride, dissolved sulphur dioxide and hydrogen chloride gases were removed by distillation of about 320ml toluene under vacuum. GC, GCMS and NMR analysis of the product were consistent with the structure of the acid chloride (IIIa). Yield, 175gm of a 54% solution of the acid chloride in toluene, ~ 97% theory.  $\alpha_D = +46^\circ$  (c= 0.012, DCM).
- 30



## EXAMPLE 2

### **Preparation of 1 R Cyhalothrin (II)**

A 2 litre jacketed non-baffled split-neck reaction flask was equipped with an overhead stirrer (turbine), thermometer, condenser, two syringe pumps and was vented to a caustic/hypochlorite scrubber. To the reactor was charged sodium cyanide (29.4g, 0.59 mol) in water (101ml) and the reaction was agitated (100 rpm) for 10 min to ensure complete dissolution of the cyanide. Hexane (79ml) was added to the reactor, then the reaction mass was cooled to 10°C and the agitation rate adjusted to 235 rpm. 1R *cis*-Z 3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethyl cyclopropanecarboxylic acid chloride (133.2g, 0.50 mol) and 3-phenoxybenzaldehyde (99g, 0.50 mol) were added simultaneously to the reactor over 3hr. Once the addition was complete the reaction was agitated for a further 17hr at 10°C. GC analysis indicated the presence of un-reacted 3-phenoxybenzaldehyde, so a further charge of 1R *cis*-Z 3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethyl cyclopropanecarboxylic acid chloride was made (3.6g, 0.014mol) and the reaction agitated for a further 2hr. The reaction mass was diluted with hexane (156ml), and the aqueous layer allowed to settle before separation. Residual cyanide in the organic phase was destroyed with sodium hypochlorite liquor to a positive show on starch/iodide paper. The positive show for hypochlorite was removed with sodium bisulphite and the aqueous phase again settled and separated. The product oil weighed 228g giving a yield of 97.8%. The composition of the product was - 1 R *cis*-Z- $\alpha$ -S diastereomer 45.5% and 1R *cis*-Z- $\alpha$ -R diastereomer 45.9%. GCMS (both diastereomers); 449 (M<sup>+</sup>), 349, 225, 208, 197, 181, 141 [α]<sub>D</sub><sup>20</sup>: +29.8° (c= 0.0104, DCM)

## EXAMPLE 3

### **Preparation of Gamma-cyhalothrin**

A 600ml vacuum jacketed non-baffled split-neck reaction flask was equipped with a dual blade agitator. To the reactor, already containing a slurry of epimerisation mass as seed from a previous experiment (already ~ 2/3rds full), was charged the 1 R cyhalothrin (II) (166.3g) in a mixture of isopropanol (99.9g), water (3.09g) and di-isopropylamine (5.22g). The reaction mass was agitated at 100rpm and -5°C for 65hr, before removing 251g of reaction mass and transferring it to a 1 litre split-neck reaction vessel in an ice/water bath. To the 1 litre reaction vessel was charged hexane (60ml) and sulphuric acid (47ml of 50% w/w), and the resultant thick paste was agitated for 0.5hr before diluting with hexane (350ml) and water (350ml). The mixture was heated to 55°C for 30min, then transferred to a separating funnel and the lower

aqueous phase was separated off. The hexane solution was concentrated under reduced pressure to give the desired product as a viscous red/brown oil which solidified upon standing to give an off-white solid. The product weighed 151.2g giving an estimated yield of 97% as gamma-cyhalothrin and a diastereomer ratio of 95:5.

5  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ): 1.20 (s, 3H, cyclopropane  $\text{CH}_3$ ), 1.30 (s, 3H, cyclopropane  $\text{CH}_3$ ), 2.00 (d, 1H,  $\text{CHCO}_2$ ), 2.25 (t, 1H,  $\text{CHCH}=\text{CClCF}_3$ ), 6.35 (s, 1H,  $\text{CHCN}$ ), 6.80 (d, 1H,  $\text{CH}=\text{CClCF}_3$ ), 7.00-7.50 (m, 9H, ArH).

GCMS; 449 ( $\text{M}^+$ ), 349, 225, 208, 197, 181, 141.

$[\alpha]_{\text{D}}^{20}$ : +36.6° (c= 0.0075, DCM)

10

#### EXAMPLE 4

##### **One-pot process for gamma-cyhalothrin**

Sodium cyanide (253 gm) and water (599ml) were charged to a 5 litre baffled reactor fitted with turbine, agitator and thermometer. The reactor contents were gently agitated to dissolve the cyanide crystals and hexane added (460ml). The contents of the reactor were then cooled to +6°C  
15 by applying coolant to the jacket and 3-phenoxy benzaldehyde (544gm) and 1R *cis*-Z 3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethyl cyclopropanecarboxylic acid chloride (736gm) were added to the reaction mass over a period of 8 hours. The resultant emulsion was then stirred overnight to complete the reaction. The solid that had formed in the reaction flask was broken down with a spatula and the agitation increased to mix the solids in. A thick slurry was produced.  
20 A small sample (~4 ml) of the slurry was filtered and the paste given a token hexane wash then pulled dry. The solid (0.7884 gm) produced was a fine white crystalline material. GC analysis of the solid showed it to be gamma-cyhalothrin with a 1R alpha-S to 1R alpha-R ratio of 86:14. (Yield approx 33%). The reaction mixture was then diluted with hexane (950ml) and the reactor temperature increased to +8°C on the reactor jacket, the reaction mass being a fairly thin slurry.  
25 The following day, a 4ml sample of reaction mass was removed and washed with hexane as before. The resultant white solid was washed with a little hexane and pulled dry. GC analysis showed that the material had a 1R alpha-S to 1R alpha-R ratio of 94:6. A total of 0.5352 gm of paste was obtained from this second filtration, which equated to a yield of approx. 32% gamma cyhalothrin. Mpt 55-58°C

30


CLAIMS

1. A process for the preparation of gamma-cyhalothrin comprising a) chlorinating 1R *cis*-Z  
5 3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethyl cyclopropanecarboxylic acid to give  
1R *cis*-Z 3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethyl cyclopropanecarboxylic  
acid chloride; b) esterifying 1R *cis*-Z 3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethyl  
cyclopropanecarboxylic acid chloride with 3-phenoxy benzaldehyde in the presence of a  
10 epimerising the diastereoisomeric mixture under conditions in which the least soluble  
diastereoisomer crystallises from solution.

2. A process according to claim 1 wherein steps b) and c) are combined in a one pot process.

15 3. A semi-batch process according to claim 1 where between 1% and 90% of the one batch  
from step c) forms the basis for nucleation and crystallization of the following batch in  
step c).

4. A compound of formula (II)



(II)